

KINECT-DCP: Clinical Trial Overview

A Phase 3, randomized, double-blind, placebo-controlled study to assess the efficacy, safety, and tolerability of valbenazine for the treatment of dyskinesia due to cerebral palsy





The information contained in these slides relates to a use of valbenazine that has not been approved by the FDA or any other regulatory agency

FDA, United States Food and Drug Administration


KINECT-DCP: Study Design^{1,2}



A Phase 3, randomized, double-blind, placebo-controlled study to assess the efficacy, safety, and tolerability of valbenazine for the treatment of dyskinesia due to cerebral palsy

~80 participants will be enrolled randomized 1:1

Valbenazine has not been approved by the FDA for the treatment of dyskinesia due to cerebral palsy



- 14-week treatment period
- Randomization 1:1 PBO to VBZ

- 32-week treatment period
- All participants receive VBZ

- ~1 year
- All OLE and ROL participants receive VBZ
- ROL:** for participants who completed the study prior to addition of the optional OLE



VBZ capsule administered once daily orally or via gastrostomy/gastrojejunostomy tube

PBO, placebo; OLE, open-label extension; ROL, re-enrolled open-label; VBZ, valbenazine 1. Clinicaltrials.gov. [NCT05206513](https://clinicaltrials.gov/ct2/show/study/NCT05206513) | 2. Data on File (VBZ-DCP-0002). Neurocrine Biosciences.

KINECT-DCP: Inclusion & Exclusion Criteria

Key Inclusion Criteria

- Male or female 6 to 70 years old
- Medically confirmed diagnosis of dyskinetic cerebral palsy (DCP): cerebral palsy with choreiform movements (i.e., a hyperkinetic movement disorder due to cerebral palsy; moderate severity or greater)
- Medical conditions are stable and expected to remain stable throughout the study

Key Exclusion Criteria

- Are pregnant or breastfeeding
- Have a clinical diagnosis or history of dyskinesia due to condition other than cerebral palsy
- Have inability to swallow soft solids, unless medications can be administered via a gastrostomy/gastrojejunostomy tube
- Have any suicidal behavior or suicidal ideation in the year prior to screening or on Day 1
- Is a substance abuser of any compound
- Known history of long QT syndrome or cardiac tachyarrhythmia, or clinically significant ECG abnormalities

KINECT-DCP: Assessments

Primary Efficacy Endpoint

- Change in the Total Maximal Chorea (**TMC**) score of the Unified Huntington's Disease Rating Scale (UHDRS) from baseline to the average of the Week 12 and Week 14 assessments

Key Secondary Endpoints

- Change in Clinical Global Impression of Severity (**CGI-S**) score from baseline to Week 14
- Change in the Movement Disorders - Childhood Rating Scale (**MD-CRS**) Part I score from baseline to the average of the Week 12 and Week 14 assessments
- Change in the Total Maximal Dystonia (**TMD**) score of the UHDRS from baseline to the average of the Week 12 and Week 14 assessments
- Patient, Caregiver, and Clinical Global Impression of Improvement (**PGI-I, CaGI-I, and CGI-I**) score at Week 14
- Goal attainment score using the Goal Attainment Scale (**GAS**) at Week 14
- Change in pain assessment from baseline to Week 14 using the Faces Pain Scale-Revised (**FPS-R**)
- Change in the UHDRS Total Motor Score (**TMS**) from baseline to the average of the Week 12 and Week 14 assessments

Safety Endpoints

- Safety assessments and monitoring will occur throughout the trial

+ UHDRS

+ CGI-S

+ MD-CRS